

Three-graviton scattering in Matrix theory revisited

Washington Taylor IV¹ and Mark Van Raamsdonk

Department of Physics
Joseph Henry Laboratories
Princeton University
Princeton, New Jersey 08544, U.S.A.
wati, mav @princeton.edu

Abstract

We consider a subset of the terms in the effective potential describing three-graviton interactions in Matrix theory and in classical eleven-dimensional supergravity. In agreement with the results of Dine and Rajaraman, we find that these terms vanish in Matrix theory. We show that the absence of these terms is compatible with the classical supergravity theory when the theory is compactified in a lightlike direction, resolving an apparent discrepancy between the two theories. A brief discussion is given of how this calculation might be generalized to compare the Matrix theory and supergravity descriptions of an arbitrary 3-body system.

June 1998

¹Address after August 1998: Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge MA 02139

In the wake of the Matrix theory conjecture of Banks, Fischler, Shenker and Susskind [1] and the AdS conjectures of Maldacena [2] there has been a remarkable body of evidence found for a deep connection between supersymmetric Yang-Mills theories and supergravity. This connection can be made very precise for certain terms arising in the perturbative one-loop expansion of Matrix theory (See [3, 4, 5] for reviews of Matrix theory). Terms in the two-graviton Matrix theory interaction potential of the form v^4/r^7 are protected by a nonrenormalization theorem [6]. These terms agree precisely with the two-graviton interaction in 11-dimensional supergravity [7, 1]. This result can be made much more general: the leading one-loop interaction potential between an arbitrary pair of Matrix theory objects is of the form F^4/r^7 , and it has been shown that this matches precisely with the potential between an arbitrary pair of objects in the linearized supergravity theory [8]. It seems likely that the full F^4/r^7 potential is protected by supersymmetry, although this has not yet been conclusively demonstrated.

At this point in time it is unclear how far the relationship extends between perturbative Matrix theory and classical supergravity. It has been shown that finite N Matrix theory corresponds to DLCQ M-theory [9, 10, 11]. This does not mean, however, that finite N Matrix theory necessarily agrees with DLCQ supergravity [3, 12]. If all terms in the perturbative Matrix theory expansion are not protected by nonrenormalization theorems, it seems necessary to understand the large N limit of the theory to verify the correspondence at higher order. It has been shown that certain higher-order terms in the Matrix theory potential agree with supergravity. Terms of the form $F^4 X^n / r^{7+n}$ precisely reproduce higher-moment interactions from supergravity [13, 8]. Some two-loop terms of the form v^6/r^{14} seem to match with supergravity [14], although the situation regarding these terms is still unclear (other terms of this form may break the correspondence [15]; there cannot, however, be v^6 terms at higher order in $1/r$ [16]). The first concrete suggestion of a breakdown of the perturbative Matrix theory/supergravity correspondence appeared in the work of Dine and Rajaraman [17]. These authors considered a 3-graviton scattering process. They found what appears to be a contradiction between perturbative Matrix theory and classical supergravity; certain terms which they expected in the classical supergravity theory could not be reproduced at any order in Matrix theory. In this letter we address this problem; for certain terms considered by Dine and Rajaraman we show that there is agreement between Matrix theory and classical null-compactified supergravity. The notation and conventions used here roughly follow those of [8].

As this letter was being written, we received the preprint [18] which addresses the same question. The results of [18] seem to be in contradiction with those presented here, although both papers reach the conclusion that Matrix theory and supergravity agree for three-graviton scattering. The results we describe here imply that the terms in the diagram computed in [18] will be canceled by terms from other diagrams, so that there is no net contribution to the terms of the form discussed in this letter. This cancellation is demonstrated explicitly in the Appendix.

The situation considered in [17] is a scattering process with three incoming and three outgoing gravitons. Gravitons 2 and 3 are separated by a distance r and gravitons 1 and 3 are separated by a distance R with $R \gg r$. Dine and Rajaraman argued that in supergravity there is a term in the amplitude proportional to

$$\frac{(v_1 - v_2)^2(v_2 - v_3)^2(v_1 - v_3)^2}{R^7 r^7} \quad (1)$$

where v_i is the velocity of the i th graviton. They showed that no such term can appear in Matrix theory, and concluded that there appears to be a discrepancy between Matrix theory and supergravity. They suggested that the resolution of this discrepancy might reside in the subtlety of the large N limit. In this letter, we argue that the discrepancy may be eliminated by taking proper account of the lightlike compactification of the supergravity theory.

In our discussion, we focus on the subset of the terms in (1) which are proportional to v_1^4 . We show that the vanishing of these terms in Matrix theory is in fact in complete agreement with what we would expect from the supergravity theory when the theory is considered in the appropriate DLCQ context. A complete demonstration of the correspondence between the $1/r^7 R^7$ terms in the perturbative supergravity potential and the Matrix theory potential for the three-graviton system would involve a more complicated calculation, and will be discussed in more detail elsewhere.

We begin by explicitly showing that there are no terms proportional to v_1^4/R^7 in the Matrix theory potential which depend on the velocities of gravitons 2 and 3. This can be demonstrated by first integrating out the off-diagonal 1-2 and 1-3 fields, and then integrating over the degrees of freedom in the 2-3 system. Considering gravitons 2 and 3 as a single system, which we denote by quantities with a hat, and graviton 1 as a second system denoted by quantities with a tilde, the leading $1/R^7$ term in the effective potential between the 2-3 system and graviton 1, which arises from integrating out the 1-2 and 1-3 fields, is given by the general two-body gravitational interaction formula [8]

$$V_{\text{one-loop}} = -\frac{15R_c^2}{4R^7} \left(\hat{\mathcal{T}}^{IJ} \tilde{\mathcal{T}}_{IJ} - \frac{1}{9} \hat{\mathcal{T}}^I{}_I \tilde{\mathcal{T}}^J{}_J \right) \quad (2)$$

where $\hat{\mathcal{T}}^{IJ}$ and $\tilde{\mathcal{T}}^{IJ}$ are defined in terms of the matrices $\hat{X}^a, \tilde{X}^a$ describing the 2-3 system and particle 1 through the expressions

$$\begin{aligned} \mathcal{T}^{--} &= \frac{1}{R_c} \text{STr} \left(\frac{1}{4} \dot{X}^a \dot{X}^a \dot{X}^b \dot{X}^b + \frac{1}{4} \dot{X}^a \dot{X}^a F^{bc} F^{bc} + \dot{X}^a \dot{X}^b F^{ac} F^{cb} \right. \\ &\quad \left. + \frac{1}{4} F^{ab} F^{bc} F^{cd} F^{da} - \frac{1}{16} F^{ab} F^{ab} F^{cd} F^{cd} \right) \\ \mathcal{T}^{-a} &= \frac{1}{R_c} \text{STr} \left(\frac{1}{2} \dot{X}^a \dot{X}^b \dot{X}^b + \frac{1}{4} \dot{X}^a F^{bc} F^{bc} + F^{ab} F^{bc} \dot{X}^c \right) \\ \mathcal{T}^{+-} &= \frac{1}{R_c} \text{STr} \left(\frac{1}{2} \dot{X}^a \dot{X}^a + \frac{1}{4} F^{ab} F^{ab} \right) \\ \mathcal{T}^{ab} &= \frac{1}{R_c} \text{STr} \left(\dot{X}^a \dot{X}^b + F^{ac} F^{cb} \right) \end{aligned}$$

$$\begin{aligned}\mathcal{T}^{+a} &= \frac{1}{R_c} \text{STr } \dot{X}^a \\ \mathcal{T}^{++} &= \frac{1}{R_c} \text{STr } \mathbb{1} = \frac{N}{R_c}\end{aligned}$$

with $F_{ab} = -i[X_a, X_b]$. (We use $I, J, \dots$ to denote 11-dimensional indices and $a, b, \dots$ to denote 9-dimensional transverse indices. R_c is the compactification radius of the x^- direction.)

For graviton 1 we have $\tilde{X}^a = (x_1^a + v_1^a t) \mathbb{1}_{N_1 \times N_1}$, so

$$\begin{aligned}\tilde{\mathcal{T}}^{++} &= \frac{N_1}{R_c} & \tilde{\mathcal{T}}^{+a} &= \frac{N_1}{R_c} v_1^a & \tilde{\mathcal{T}}^{ab} &= \frac{N_1}{R_c} v_1^a v_1^b \\ \tilde{\mathcal{T}}^{+-} &= \frac{N_1}{2R_c} v_1^2 & \tilde{\mathcal{T}}^{-a} &= \frac{N_1}{2R_c} v_1^a v_1^2 & \tilde{\mathcal{T}}^{--} &= \frac{N_1}{4R_c} v_1^4\end{aligned}\tag{3}$$

If we restrict attention to the terms in the effective potential which are proportional to v_1^4/R^7 , we are left with the expression

$$-\frac{15R_c N_1}{16R^7} \langle \hat{\mathcal{T}}^{++} \rangle v_1^4$$

where by $\langle \cdot \rangle$ we mean the expectation value after integration over the fluctuations of the 2-3 system. Since $\hat{\mathcal{T}}^{++} = (N_2 + N_3)/R_c$ is a constant, however, we see that there are no corrections at any order in $1/r$ to the one-loop term in the interaction potential

$$-\frac{15N_1(N_2 + N_3)}{16R^7} v_1^4.\tag{4}$$

This agrees with the result of Dine and Rajaraman in [17] that no terms of the form (1) can be generated in Matrix theory after summing all diagrams at any loop order.

We now discuss the corresponding terms in the classical supergravity potential. The leading $1/R^7$ term in the classical supergravity potential between the 2-3 system and particle 1 arises from diagrams in which a single graviton is exchanged. Because the propagator for a graviton with zero longitudinal momentum contains a delta function $\delta(x^+ - y^+)$, the resulting interaction is a classical instantaneous effective potential which takes precisely the form (2), where $\hat{\mathcal{T}}^{IJ}$ and $\tilde{\mathcal{T}}^{IJ}$ are the integrated components of the classical stress tensor for the 2-3 and 1 systems. Note that the stress tensor $\hat{\mathcal{T}}$ in general contains terms proportional to $1/r^7$ which arise from internal graviton exchange processes between particles 2 and 3. Such terms correspond in the classical theory to nonlinear corrections to the stress tensor from the gravitational field itself. Since particle 1 is pointlike, its classical stress tensor components are given by

$$\tilde{\mathcal{T}}^{IJ} = \frac{p^I p^J}{p^+}.$$

Using $p^+ = N_1/R_c$, $p^a = p^+ v_1^a$, and $p^- = p_a^2/2p^+$ we arrive at the expressions (3) for the components of the integrated stress tensor of particle 1. To determine the complete set of terms in the classical supergravity potential which are proportional to v_1^4/R^7 it remains to evaluate the component $\hat{\mathcal{T}}^{++}$ of the integrated stress-tensor for the 2-3 system. This,

however, is precisely the total conserved momentum of the system in the x^- direction, which in the null-compactified supergravity theory corresponds to the quantity $(N_2 + N_3)/R_c$. This shows that the only terms in the DLCQ supergravity three-graviton potential proportional to v_1^4/R^7 are contained in the linearized supergravity interaction (4), which is precisely reproduced by Matrix theory.

We have therefore shown that Matrix theory and DLCQ supergravity are in complete agreement for a subset of the terms in the three-graviton potential. This resolves, at least in part, the discrepancy found by Dine and Rajaraman. At this point, a few comments may be helpful in clarifying the relationship between the analysis above and that in [17]. In [17], the authors computed a term proportional to (1). This term is also proportional to the product of the momentum components in the x^- direction

$$k_1^+ k_2^+ k_3^+ \frac{(v_1 - v_2)^2 (v_2 - v_3)^2 (v_1 - v_3)^2}{R^7 r^7}. \quad (5)$$

In [17] there was an implicit assumption that $k_i^+ = N_i/R_c$ for $i = 1, 2, 3$. However, this identification is not necessarily correct. Gravity is a nonlinear theory. If we think of supergravity as a field theory with a spin 2 field h_{IJ} living in a flat space-time background η_{IJ} , so that $g_{IJ} = \eta_{IJ} + h_{IJ}$, then all components of the stress tensor, including the momentum $\mathcal{T}^{++}$, contain contributions nonlinear in the field h_{IJ} . The total momentum of the 2-3 system is then given by $\hat{\mathcal{T}}^{++} = k_2^+ + k_3^+ + \text{nonlinear terms}$. These nonlinear terms may be explicitly computed by expanding the stress tensor components order by order in h (see for example [19]). In the case of interest here, it can be verified that there are nonlinear terms of the form $k_2^+ k_3^+ (v_2 - v_3)^2 / r^7$ which contribute to the total $\hat{\mathcal{T}}^{++}$. There are numerous subtleties involved with defining DLCQ theories (see for example [12]). Nonetheless, it seems reasonable to expect that the classical limit of DLCQ supergravity will simply be the classical supergravity theory compactified on a null circle. In this classical theory, the discretized quantity $\hat{\mathcal{T}}^{++}$ is the *total* momentum in the compact direction, and includes all terms nonlinear in h . It follows that the nonlinear terms (5) are automatically included in the result (4) when $(N_2 + N_3)/R_c$ is identified with the total $\hat{\mathcal{T}}^{++}$. This explains in detail how the apparent mystery of the missing terms is resolved.

We conclude with a brief discussion of how the analysis in this letter might be generalized. For an arbitrary 3-body system, and for the 3-graviton system in particular, there are terms in the supergravity potential proportional to

$$\frac{\hat{\mathcal{T}}^{(2)} \tilde{\mathcal{T}}}{R^7 r^7} \quad (6)$$

where $\hat{\mathcal{T}}^{(2)}$ represents a contribution to the total stress tensor of the 2-3 system arising from quadratic terms in the gravity theory. For general objects in the supergravity theory there will also be membrane current interactions arising from 3-form exchange. In general, we would expect to be able to reproduce terms of the form (6) by integrating out the 2-3 fields in the expression (2). In the case where we restrict attention to terms proportional to the

fourth power of the curvature $F^4(v^4)$ of object 1, the integral over the 2-3 fields is trivial because the integrand is a constant. This is the case we have considered in this letter. For the other terms, the calculation is more complicated. A complete calculation of $\langle \hat{\mathcal{T}}^{IJ} \rangle$ would require, in particular, a generalization of the results of [8] describing the components of the stress tensor in the presence of arbitrary background fermion fields. Work in this direction is in progress [20].

It was recently shown by Paban, Sethi and Stern [16] that the v^6 terms in the two-graviton interaction potential are not renormalized. This gives some hope that the full collection of terms of the form F^6/r^{14} , including those relevant for the general 3-body calculation just mentioned, may similarly be protected by supersymmetry. If true, this would indicate that Matrix theory correctly reproduces supergravity to quadratic order. Recent work [21] indicates that there may be problems with the velocity expansion at order v^8 and beyond. This suggests that the correspondence with supergravity may break down at cubic order; more detailed calculations are probably necessary, however, to be certain of this conclusion.

Note added: Soon after the preprint of this paper appeared on the net, a number of related papers appeared [22, 23, 24, 25] discussing the three-graviton issue in Matrix theory. We have added this short note to clarify how this paper is related to these other papers. The paper [23] contains an analysis of the cancellation of the diagram considered in [18]. The result of [23] agrees with the result of the Appendix of this letter, which states that in the background field gauge we use here, this diagram is exactly canceled at one-loop order. In [25] it is suggested that this particular diagram might not cancel at one-loop order in a different gauge. After summing all diagrams, however, it is clear that physical quantities such as the phase shift in the scattering amplitude should be gauge-invariant, so one would expect terms of the form discussed in this paper to cancel after summing all diagrams even in the gauge of [25]. The comprehensive analysis of Okawa and Yoneya [22] makes it clear that there is an exact agreement between Matrix theory and supergravity for the three-graviton scattering process (at least when recoil is neglected). In particular, by taking the limit of their expression (4.4) in the case of three particles A, B and C where $r = x_{BC} \ll R = x_{AC}$ it can be seen that no terms proportional to $v_A^4/r^7 R^7$ appear. The essential result of this letter is a simple argument for the vanishing of this term both in Matrix theory and supergravity. The vanishing of this term in the supergravity theory indicates that the analysis of Dine and Rajaraman in [17] was incomplete. It should be noted that the physical argument given in this letter for the vanishing of this term in supergravity does not depend on the null quantization of the gravity theory; in fact, the identification of $(N_2 + N_3)/R_c$ with the integrated T^{++} of the 2-3 system can be seen simply from the fact that this is a conserved quantity in both the Matrix and supergravity theories, which can be identified in this way when particles 2 and 3 are widely separated, and which therefore can also be identified as the particles come together. Thus, regardless of whether the supergravity is null-compactified or not, the nonlinear terms we have discussed in T^{++} (these terms are also discussed in

[24]) must automatically be included in $(N_2 + N_3)/R_c$. In summary, the results of this letter give a simple physical argument indicating that the results of Dine and Rajaraman are incomplete. The result presented here agrees with the paper of Okawa and Yoneya [22], in which the question is settled decisively by a careful calculation of the three-graviton scattering amplitude in both theories. The methods used in this paper can be naturally extended to study interactions between n arbitrary Matrix theory objects, and to address the question of the Matrix theory/supergravity correspondence at higher order.

Acknowledgments

We would like to thank T. Banks, M. Dine, D. Kabat, P. Kraus, D. Lowe and V. Periwal for helpful conversations and correspondence. The work of MVR is supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC). The work of WT was supported in part by the National Science Foundation (NSF) under contracts PHY96-00258 and PHY94-07194. WT would like to thank the organizers of the Duality 98 program and the ITP in Santa Barbara for hospitality while this work was being completed.

Appendix

In this appendix we describe explicitly the cancellation of the term calculated in [18]². In that paper a single diagram, the “setting sun” diagram, is computed. This computation is performed in two steps: first the very massive 1-2 and 1-3 fields are integrated out, then the 2-3 fields are integrated out. The cancellation of the setting-sun diagram occurs already after integrating out the very massive modes, when the corresponding setting-sun diagram is computed with fermionic 1-2 and 1-3 fields. This conclusion essentially follows from the computation in [26], where the one-loop calculation was performed for an arbitrary bosonic background. The connection between these calculations is somewhat subtle, however, as they are performed using different propagators for the off-diagonal fields.

In order to make the cancellation mechanism completely clear in the case discussed in [18], we describe here explicitly the diagrammatic cancellation using the same propagators as those used in that paper. We make the discussion slightly more general by considering n gravitons at positions R_i where $R_1 \gg R_i$ for $i > 1$. We assume that the $(N - 1) \times (N - 1)$ background matrices describing gravitons $2 - n$ contain arbitrary (but small) off-diagonal terms, and write these matrices as

$$X_{ij}^a = R_{ij}^a + K_{ij}^a, \quad 1 < i, j \leq n$$

where $R^a = \text{Diag}(R_2^a, \dots, R_n^a)$. We wish to integrate out the off-diagonal bosonic and fermionic fields x_i^a ($1 \leq a \leq 9$) and ψ_i^α ($1 \leq \alpha \leq 16$) (where we have simplified nota-

²After this work was completed we were informed that this cancellation has been independently derived by Echols and Gray [23]

tion by writing, for example, $x_i = x_{1i}$). We ignore the gauge fluctuations in this calculation as they are not relevant to terms of the type computed in [18]. The relevant terms in the Lagrangian are

$$\frac{1}{R_c} \left\{ \dot{x}_i^* \dot{x}_i - x_i^* (r_i^2) x_i + x_i^* (r_i \cdot K_{ij} + K_{ij} \cdot r_j - (K \cdot K)_{ij}) x_j - 2(x_i^a)^* ([Y_a, Y_b])_{ij} x_j^b \right. \\ \left. + i\psi_i \dot{\psi}_i + \psi_i \gamma^a r_i^a \psi_i - \psi_i \gamma^a (K^a)_{ij} \psi_j \right\}$$

where we have defined

$$(Y^a)_{ij} = (X^a)_{ij} - R_1^a \delta_{ij}$$

and

$$r_i = R_1 - R_i,$$

and we have suppressed some spatial and spinor indices a and α . Treating r_i as the masses of the bosonic and fermion fields, and the remaining terms as interactions, we can perform the one-loop calculation order by order by summing over bosonic diagrams with insertions of the quadratic vertices of the forms $r \cdot K$, K^2 and $[Y, Y]$, and fermionic diagrams with insertions of the quadratic vertex of the form $K \cdot \gamma$. Terms with fewer than four bosonic vertices of the $[Y, Y]$ type are all canceled, as shown in [26]. In particular, the diagram calculated in [18] comes from a term of the form

$$2\langle ((x_i^a)^* ([Y^a, Y^b])_{ij} x_j^b) \cdot ((x_k^c)^* ([Y^c, Y^d])_{kl} x_l^d) \rangle.$$

Using the leading part of the bosonic propagator [27, 18]

$$\Delta(t_1, t_2 | r^2) = \frac{1}{2r} e^{-r|t_1 - t_2|}$$

we find that this term gives a contribution of

$$\frac{2}{r_i r_j (r_i + r_j)} \left\{ ([R^a, K^b])_{ij} ([R^b, K^a])_{ji} - ([R^a, K^b])_{ij} ([R^a, K^b])_{ji} \right\} \quad (7)$$

as well as terms of the form $[K, K]^2$. The expression (3.12) in [18] is a piece of the term (7) in the case $n = 2$.

Now consider the fermionic loop diagram with two insertions of $K^a \gamma^a$. The leading fermionic propagator contains two terms, one containing a θ function and no γ matrices, and the other given by

$$\frac{r^a \gamma^a}{2r} e^{-r|t_1 - t_2|}.$$

Using only this part of the propagator, we get a contribution

$$-\frac{1}{4r_i r_j (r_i + r_j)} \text{Tr} (\gamma^a \gamma^b \gamma^c \gamma^d r_i^a K_{ij}^b r_j^c K_{ji}^d)$$

The trace can be rewritten in the form

$$\begin{aligned}
\text{Tr}(\cdot) &= 16 \text{Tr} \left(r^a K^b r^b K^a - r^a K^b r^a K^b + r^a K^a r^b K^b \right) \\
&= 8 \left(([R^a, K^b])_{ij} ([R^b, K^a])_{ji} - ([R^a, K^b])_{ij} ([R^a, K^b])_{ji} \right) \\
&\quad + 8 \left((r \cdot K + K \cdot r)_{ij} (r \cdot K + K \cdot r)_{ji} \right) - 16 r_j^2 (K \cdot K)_{jj}
\end{aligned}$$

The first term in this expression precisely cancels (7). The remaining terms are higher moments of canceling diagrams, and are canceled by other one-loop diagrams. The second term is canceled by the bosonic diagram with two $(r \cdot K)$ insertions, and the third term is canceled by a combination of the bosonic term with a single $K \cdot K$ term and the fermionic diagram with two $K \cdot \gamma$ insertions and theta functions in the propagators.

We have thus shown explicitly, using the same propagator structure as in [18], that in general all the bosonic one-loop terms with two $[Y, Y]$ insertions are canceled by a combination of bosonic and fermionic diagrams. In particular, this shows explicitly that the term found in expression (3.12) of [18] is canceled simply by integrating out the very massive 1-2 and 1-3 modes, in accord with the results of Dine and Rajaraman and the discussion in the main text of this letter.

References

- [1] T. Banks, W. Fischler, S. Shenker, and L. Susskind, “M Theory as a Matrix Model: A Conjecture,” *Phys. Rev.* **D55** (1997) 5112, [hep-th/9610043](#).
- [2] J. Maldacena, “The large N limit of superconformal field theories and supergravity,” [hep-th/9711200](#).
- [3] T. Banks, “Matrix Theory,” *Nucl. Phys. Proc. Suppl.* **67** (1998) 180, [hep-th/9710231](#).
- [4] D. Bigatti and L. Susskind, “Review of matrix theory,” [hep-th/9712072](#).
- [5] W. Taylor, “Lectures on D-branes, gauge theory and M(atrices),” Proceedings of Trieste summer school 1997, to appear; [hep-th/9801182](#).
- [6] S. Paban, S. Sethi and M. Stern, “Constraints from extended supersymmetry in quantum mechanics,” [hep-th/9805018](#).
- [7] M. R. Douglas, D. Kabat, P. Pouliot, and S. Shenker, “D-Branes and Short Distances in String Theory,” *Nucl. Phys.* **B485** (1997) 85, [hep-th/9608024](#).
- [8] D. Kabat and W. Taylor, “Linearized supergravity from Matrix theory,” *Phys. Lett.* **B**, to appear; [hep-th/9712185](#).
- [9] L. Susskind, “Another Conjecture about M(atrrix) Theory,” [hep-th/9704080](#).

- [10] A. Sen, “D0 Branes on T^n and Matrix Theory,” [hep-th/9709220](#).
- [11] N. Seiberg, “Why is the Matrix Model Correct?,” *Phys. Rev. Lett.* **79** (1997) 3577, [hep-th/9710009](#).
- [12] S. Hellerman and J. Polchinski, “Compactification in the Lightlike Limit,” [hep-th/9711037](#).
- [13] W. Taylor and M. Van Raamsdonk, “Angular momentum and long-range gravitational interactions in Matrix theory,” [hep-th/9712159](#).
- [14] K. Becker, M. Becker, J. Polchinski, and A. Tseytlin, “Higher Order Graviton Scattering in M(atrix) Theory,” *Phys. Rev.* **D56** (1997) 3174, [hep-th/9706072](#).
- [15] V. Periwal and R. von Unge, “Accelerating D-branes,” [hep-th/9801121](#).
- [16] S. Paban, S. Sethi and M. Stern, “Supersymmetry and higher-derivative terms in the effective action of Yang-Mills theories,” [hep-th/9806028](#).
- [17] M. Dine and A. Rajaraman, “Multigraviton Scattering in the Matrix Model,” [hep-th/9710174](#).
- [18] M. Fabbrichesi, G. Ferretti and R. Iengo “Supergravity and matrix theory do not disagree on multi-graviton scattering,” [hep-th/9806018](#).
- [19] S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972).
- [20] P. Kraus, W. Taylor and M. Van Raamsdonk, work in progress.
- [21] M. Dine, R. Echols and J. P. Gray, “Renormalization of higher derivative operators in the matrix model,” [hep-th/9805007](#).
- [22] Y. Okawa and T. Yoneya, “Multi-body interactions of D-particles in supergravity and Matrix theory,” [hep-th/9806108](#).
- [23] R. Echols and J. P. Gray, “Comment on multigraviton scattering in the matrix model,” [hep-th/9806109](#).
- [24] J. McCarthy, L. Susskind and A. Wilkins, “Large N and the Dine-Rajaraman problem”, [hep-th/9806136](#)
- [25] M. Fabbrichesi, G. Ferretti and R. Iengo “About the Matrix model computation of three-graviton scattering,” [hep-th/9806166](#).
- [26] D. Kabat and W. Taylor, “Spherical membranes in Matrix theory,” *Adv. Theor. Math. Phys.* **2**, 181-206, [hep-th/9711078](#).

- [27] K. Becker and M. Becker, “A Two-Loop Test of M(atrix) Theory,” *Nucl. Phys.* **B506** (1997) 48-60, [hep-th/9705091](#).